arXiv:math/0410322v1 [math.QA] 13 Oct 2004

New approach to Hermitian q-differential operators on $\mathbb{R}_q^N$

Gaetano Fiore *

Dip. di Matematica e Applicazioni, Fac. di Ingegneria
Università di Napoli, V. Claudio 21, 80125 Napoli
and
I.N.F.N., Sezione di Napoli,
Complesso MSA, V. Cintia, 80126 Napoli

Abstract

We report on our recent breakthrough [9] in the costruction for $q>0$ of Hermitean and "tractable" differential operators out of the $U_qso(N)$-covariant differential calculus on the noncommutative manifolds $\mathbb{R}_q^N$ (the socalled "quantum Euclidean spaces").

1 Introduction

Noncommutative geometry for non-compact noncommutative manifolds is usually much more difficult than for compact ones, especially when trying to proceed from an algebraic to a functional-analytical treatment (see e.g. [12]). Major sources of complications are $\star$-structures and $\star$-representations of the involved algebras. As a noncommutative space here we adopt the socalled N-dimensional quantum Euclidean space $\mathbb{R}_q^N$ ($N \geq 3$), a deformation of $\mathbb{R}^N$ characterized by its covariance under the quantum group $U_qso(N)$ (with q real so that the latter is compact; for $\star$-representation purposes we shall later need and choose $q>0$) and shall denote by F the $\star$-algebra "of functions on $\mathbb{R}_q^N$". The $\star$-structure [13] associated to the corresponding $U_qso(N)$-covariant differential calculus [1]

---

*Talk given at the Conference "Noncommutative Geometry and Representation Theory in Mathematical Physics", Karlstad (Sweden), July 2004. Preprint 04-45 Dip. Matematica e Applicazioni, Università di Napoli; DSF/04-35 ; ESI 1529. Work partially supported by the European Commission RTN Programme HPRN-CT-2000-00131 and by MIUR

on $\mathbb{R}_q^N$ [6] is however characterized by an unpleasant nonlinear action on the differentials, the partial derivatives and the exterior derivative. This at the origin of a host of formal and substantial complications: it seems hardly possible to construct physically relevant differential operators (e.g. momentum components, Laplacian, kinetic term in a would-be field theory on $\mathbb{R}_q^N$) which are at the same time Hermitean and "tractable".

Here we report on a recent breakthrough [9] to the approach to these problems. This is based on our result (Thm 1 in [9]) that the transformation laws [13] of the N partial derivatives ∂^α and of the exterior derivative d of the differential calculus [1] on $\mathbb{R}_q^N$ under the $\star$-structure can be expressed as the similarity transformations

$$\partial^{\alpha\star} = -\tilde{\nu}'^{-2}\tilde{\partial}^\alpha\tilde{\nu}'^2, \qquad\qquad d^\star = -\tilde{\nu}'^2 d\tilde{\nu}'^{-2}, \tag{1}$$

where $\tilde{\nu}'$ is a $U_q so(N)$-invariant positive-definite pseudodifferential operator, more precisely the realization of the fourth root of the ribbon element of the extension of $U_q so(N)$ with a central element generating dilatations of $\mathbb{R}_q^N$.

The situation reminds us of what one encounters in functional analysis on the real line when taking the Hermitean conjugate of a differential operator like

$$\partial := \tilde{\nu}'(x)\,\frac{d}{dx}\,\frac{1}{\tilde{\nu}'(x)},$$

where $\tilde{\nu}'(x)$ is a positive smooth function; as an element of the Heisenberg algebra ∂ is not imaginary (excluding the trivial case $\tilde{\nu}' \equiv 1$) w.r.t. the $\star$-structure

$$x^\star = x, \qquad \left(\frac{d}{dx}\right)^\star = -\frac{d}{dx},$$

but fulfills the similarity transformation (1). This corresponds to the fact that it is not antihermitean as an operator on $L^2(\mathbb{R})$. ∂ is however (formally) antihermitean on $L^2(\mathbb{R}, \tilde{\nu}'^{-2}dx)$, the Hilbert space of square-integrable functions over $\mathbb{R}$ endowed with measure $\tilde{\nu}'^{-2}dx$. In other words, if we introduce the scalar product

$$(\boldsymbol{\phi}, \boldsymbol{\psi}) = \int \boldsymbol{\phi}^\star(x)\,\tilde{\nu}'^{-2}(x)\,\boldsymbol{\psi}(x)dx, \tag{2}$$

[as one does when setting the Sturm-Liouville problem for ∂^2], ∂ becomes antihermitean under the corresponding Hermitean conjugation $\dagger$[1]:

$$(A^\dagger\boldsymbol{\phi}, \boldsymbol{\psi}) := (\boldsymbol{\phi}, A\boldsymbol{\psi}) \qquad \Rightarrow \qquad \partial^\dagger = -\partial.$$

The idea is to adopt the analog of (2), where now ν' is a *pseudodifferential operator* rather than a function, as a scalar product also in our noncommutative space.

---

[1] The Hermitean conjugation $\dagger$ is the representation of the following modified $\star$-structure $\star'$ of the Heisenberg algebra: $a^{\star'} = (\tilde{\nu}'^{-2}\,a\tilde{\nu}'^2)^\star = \tilde{\nu}'^2\,a^*\tilde{\nu}'^{-2}$.

2 Preliminaries and notation

The algebras.

F is essentially the unital associative algebra over $\mathbb{C}[[h]]$ generated by N elements x^α (the cartesian "coordinates") modulo the relations (3) given below, and is extended to include formal power series in the generators; out of F one can extract subspaces consisting of elements that can be considered integrable or square-integrable functions. The $U_qso(N)$-covariant differential calculus on $\mathbb{R}_q^N$ [1] is defined introducing the invariant exterior derivative d, satisfying nilpotency and the Leibniz rule $d(fg) = dfg + fdg$, and imposing the covariant commutation relations (4) between the x^α and the differentials $\xi^\alpha := dx^\alpha$. Partial derivatives are introduced through the decomposition $d =: \xi^\alpha \partial_\alpha$. All the other commutation relations are derived by consistency. The complete list is given by

$$\mathcal{P}_a{}^{\alpha\beta}_{\gamma\delta} x^\gamma x^\delta = 0, \tag{3}$$

$$x^\gamma \xi^\alpha = q\hat{R}^{\gamma\alpha}_{\beta\delta} \xi^\beta x^\delta, \tag{4}$$

$$(\mathcal{P}_s + \mathcal{P}_t)^{\alpha\beta}_{\gamma\delta} \xi^\gamma \xi^\delta = 0, \tag{5}$$

$$\mathcal{P}_a{}^{\alpha\beta}_{\gamma\delta} \partial_\beta \partial_\alpha = 0, \tag{6}$$

$$\partial_\alpha x^\beta = \delta^\beta_\alpha + q\hat{R}^{\beta\gamma}_{\alpha\delta} x^\delta \partial_\gamma, \tag{7}$$

$$\partial^\gamma \xi^\alpha = q^{-1}\hat{R}^{\gamma\alpha}_{\beta\delta} \xi^\beta \partial^\delta. \tag{8}$$

Here the $N^2 \times N^2$ matrix $\hat{R}$ is the braid matrix of $SO_q(N)$ [6]. The matrices $\mathcal{P}_s$, $\mathcal{P}_a$, $\mathcal{P}_t$ are $SO_q(N)$-covariant deformations of the symmetric trace-free, antisymmetric and trace projectors respectively, which appear in the orthogonal projector decomposition of $\hat{R}$

$$\hat{R} = q\mathcal{P}_s - q^{-1}\mathcal{P}_a + q^{1-N}\mathcal{P}_t. \tag{9}$$

Thus they satisfy the equations (with $A, B = s, a, t$)

$$\mathcal{P}_A \mathcal{P}_B = \mathcal{P}_A \delta_{AB}, \qquad \mathcal{P}_s + \mathcal{P}_a + \mathcal{P}_t = \mathbf{1}_{N^2}. \tag{10}$$

The $\mathcal{P}_t$ projects on a one-dimensional sub-space and can be written in the form

$$\mathcal{P}_t{}^{\alpha\beta}_{\gamma\delta} \sim g^{\alpha\beta} g_{\gamma\delta}; \tag{11}$$

here the $N \times N$ matrix $g_{\alpha\beta}$ is a $SO_q(N)$-isotropic tensor, deformation of the ordinary Euclidean metric. The elements

$$r^2 \equiv x \cdot x := x^\alpha g_{\alpha\beta} x^\beta, \qquad \partial \cdot \partial := g^{\alpha\beta} \partial_\beta \partial_\alpha$$

are $U_qso(N)$-invariant and respectively generate the centers of F, F'. r^2 is real and positive-definite w.r.t. the $\star$-structure in F for real q introduced in [6]. $\partial \cdot \partial$ is a deformation of the Laplacian on $\mathbb{R}^N$. We shall adopt here mainly a set $\{x^\alpha\}$ of *real* generators $x^\alpha = V^\alpha_i x^i$, whereas by $\{x^i\}$ we mean the standard non-real basis adopted since the works [6, 1], in which r^2 takes the form $r^2 = \sum_i x^{i\star} x^i$ and the relations (3-8) take the simplest explicit form, and V is the (complex) linear transformation relating the two bases.[2]

We shall slightly extend F by introducing the square root r of r^2 and its inverse r^{-1} as new (central) generators; r is "the deformed Euclidean distance of the generic point of coordinates (x^i) of $\mathbb{R}^N_q$ from the origin". The elements

$$t^\alpha := r^{-1} x^\alpha$$

fulfill $t{\cdot}t = 1$ and thus generate the subalgebra $F(S^{N-1}_q)$ of "functions on the unit quantum Euclidean sphere". $F(S^{N-1}_q)$ can be completely decomposed into eigenspaces V_l of the quadratic Casimir of $U_q so(N)$, or equivalently of the Casimir w defined in (2) with eigenvalues $w_l := q^{-l(l+N-2)}$, implying a corresponding decomposition for F:

$$F(S^{N-1}_q) = \bigoplus_{l=0}^{\infty} V_l, \qquad\qquad F = \bigoplus_{l=0}^{\infty} \left(V_l \otimes \mathbb{C}[[r, r^{-1}]]\right). \tag{12}$$

An orthonormal basis $\{S^I_l\}$ (consisting of 'spherical harmonics') of V_l can be extracted out of the set of homogeneous, completely symmetric and trace-free polynomials of degree l, suitably normalized. Therefore for the generic $f \in F$

$$f = \sum_{l=0}^{\infty} f_l = \sum_{l=0}^{\infty} \sum_I S^I_l f_{l,I}(r). \tag{13}$$

We shall call $\mathcal{DC}^*$ (differential calculus algebra on $\mathbb{R}^N_q$) the unital associative algebra over $\mathbb{C}[[h]]$ generated by $x^\alpha, \xi^\alpha, \partial_\alpha$ modulo relations (3-8). We shall denote by $\bigwedge^*$ (exterior algebra, or algebra of exterior forms) the graded unital subalgebra generated by the ξ^α alone, with grading $\natural \equiv$the degree in ξ^α, and by $\bigwedge^p$ (vector space of exterior p-forms) the component with grading $\natural = p$, $p = 0, 1, 2....$ Each $\bigwedge^p$ carries an irreducible representation of $U_q so(N)$, and its dimension is the binomial coefficient $\binom{N}{p}$ [10], exactly as in the $q = 1$ (i.e. undeformed) case. We shall endow $\mathcal{DC}^*$ with the same grading $\natural$, and call $\mathcal{DC}^p$ its component with grading $\natural = p$. The elements of $\mathcal{DC}^p$ can be considered differential-operator-valued p-forms. We

---

[2] For instance, relations (3), (5) for $\mathbb{R}^3_q$ in the basis $\{x^i\} = \{x^-, x^0, x^+\}$ [6, 1] amount to:

$$x^- x^0 = q x^0 x^- \qquad x^0 x^+ = q x^+ x^0, \qquad [x^-, x^+] = (q^{\frac{1}{2}} - q^{-\frac{1}{2}}) x^0 x^0,$$

$$q\xi^- \xi^0 + \xi^0 \xi^- = q\xi^0 \xi^+ + \xi^+ \xi^0 = \xi^- \xi^+ + \xi^+ \xi^- = (\xi^\pm)^2 = (\xi^0)^2 - (q^{\frac{1}{2}} - q^{-\frac{1}{2}}) \xi^- \xi^+ = 0.$$

The $\star$-structure acts on the coordinates as follows:

$$(x^-)^* = q^{\frac{1}{2}} x^+, \qquad (x^0)^* = x^0, \qquad (x^+)^* = q^{-\frac{1}{2}} x^-.$$

shall denote by $\mathcal{H}$ (Heisenberg algebra on $\mathbb{R}_q^N$) the unital subalgebra generated by the $x^\alpha, \partial_\alpha$. Note that by definition $\mathcal{DC}^0 = \mathcal{H}$, and that both $\mathcal{DC}^*$ and $\mathcal{DC}^p$ are $\mathcal{H}$-bimodules. We shall denote by Ω^* (algebra of differential forms) the graded unital subalgebra generated by the ξ^α, x^α, with grading $\natural$, and by Ω^p (space of differential p-forms) its component with grading p; by definition $\Omega^0 = F$ itself. Clearly both Ω^* and Ω^p are F-bimodules. Finally, we recall that by replacing $q \to q^{-1}, \hat{R} \to \hat{R}^{-1}$ in (3-8) one gets an alternative $U_qso(N)$-covariant calculus, whose exterior/partial derivative and differentials $\hat{d}, i\hat{\partial}^\alpha, \hat{\xi}^\alpha$ are the $\star$-conjugates of $d, i\partial^\alpha, \xi^\alpha$ and are related to the latter by a rather complicated rational transformation [14].

Hodge duality.

The "Hodge map" [10] is a $U_qso(N)$-covariant, $\mathcal{H}$-bilinear map

$$* : \mathcal{DC}^p \to \mathcal{DC}^{N-p} \tag{14}$$

($p = 0, 1, ..., N$) such that ${}^*\mathbf{1} = dV$ and on each $\mathcal{DC}^p$ (hence on the whole $\mathcal{DC}^*$)

$$*^2 \equiv * \circ * = \mathrm{id}\,, \tag{15}$$

defined by setting on the monomials in the ξ^α

$${}^*(\xi^{\alpha_1}...\xi^{\alpha_p}) = q^{-N(p-N/2)} c_p\, \xi^{\alpha_{p+1}}...\xi^{\alpha_N}\, \varepsilon_{\alpha_N...\alpha_{p+1}}{}^{\alpha_1...\alpha_p} \Lambda^{2p-N}. \tag{16}$$

Here c_p is a suitable normalization factor [going to $1/(N-p)!$ in the commutative limit $q \to 1$], $\varepsilon^{\alpha_1...\alpha_N}$ is the q-deformed ε-tensor and $\Lambda^{\mp 1}$ are the square root and the inverse square root of

$$\Lambda^{-2} := 1 + (q^2-1)x^\alpha\partial_\alpha + \frac{q^{N-2}(q^2-1)^2}{(1+q^{N-2})^2} r^2 \partial\cdot\partial \equiv 1 + O(h); \tag{17}$$

as a pseudodifferential operator Λ acts as a dilatation:

$$\Lambda x^\alpha = q^{-1}x^\alpha\Lambda, \qquad \Lambda\partial^\alpha = q\partial^\alpha\Lambda, \qquad \Lambda\xi^\alpha = \xi^\alpha\Lambda, \qquad \Lambda 1 = 1. \tag{18}$$

In $\mathcal{DC}^*$ one can introduce [9, 2] as an alternative basis of 1-forms a "frame" [4] $\{\theta^\alpha\}$, in the sense that $[\theta^\alpha, \mathcal{H}] = 0$ (implying in particular $[\theta^\alpha, F] = 0$). In the latter (16) takes the form

$${}^*(\theta^{\alpha_1}\theta^{\alpha_2}...\theta^{\alpha_p}) = c_p\, \theta^{\alpha_{p+1}}...\theta^{\alpha_N}\, \varepsilon_{\alpha_N...\alpha_{p+1}}{}^{\alpha_1...\alpha_p}, \tag{19}$$

Restricting the domain of $*$ to the unital subalgebra $\widetilde{\Omega}^* \subset \mathcal{DC}^*$ generated by $x^\alpha, \xi^\alpha, \Lambda^{\pm 1}$ one obtains a $U_qso(N)$-covariant, $\widetilde{F}$-bilinear map (with $\widetilde{F} \equiv \widetilde{\Omega}^0$)

$$* : \widetilde{\Omega}^p \to \widetilde{\Omega}^{N-p} \tag{20}$$

fulfilling again ${}^*\mathbf{1} = dV$ and (15). Finally, introducing the exterior coderivatives

$$\delta := -{}^*\, d\, {}^* \qquad\qquad \hat{\delta} := -{}^*\, \hat{d}\, {}^* \tag{21}$$

one finds that on all of $\mathcal{DC}^*$, and in particular on all of Ω^*, the Laplacians

$$\hat{\Delta} := d\,\delta + \delta\,d = -q^2\,\partial\cdot\partial\Lambda^2 = -q^{-N}\hat{\partial}\cdot\hat{\partial}, \tag{22}$$
$$\Delta := \hat{d}\,\hat{\delta} + \hat{\delta}\,\hat{d} = -q^{-2}\,\hat{\partial}\cdot\hat{\partial}\Lambda^{-2} = -q^{N}\,\partial\cdot\partial \tag{23}$$

are respectively quadratic in the $\hat{\partial}_\alpha, \partial_\alpha$ (note: not in the $\partial_\alpha, \hat{\partial}_\alpha$!).

The restriction (20) is the notion closest to the conventional notion of a Hodge map on $\mathbb{R}_q^N$: as a matter of fact, there is no F-bilinear restriction of $*$ to Ω^*. However $\widetilde{\Omega}^*$ is not closed under the $\star$-structure which we shall recall below.

$\widetilde{U_qso(N)}$ and its realization by (pseudo)differential operators.

We extend [11] the Hopf algebra $U_qso(N)$ by adding a central, primitive generator η

$$\phi(\eta) = \mathbf{1}\otimes\eta + \eta\otimes\mathbf{1}, \qquad \epsilon(\eta) = 0, \qquad S\eta = -\eta$$

(here ϕ, ϵ, S respectively denote the coproduct, counit, antipode), and we endow the resulting Hopf algebra $\widetilde{U_qso(N)}$ by the quasitriangular structure

$$\tilde{\mathcal{R}} := \mathcal{R}\ q^{\eta\otimes\eta}, \tag{24}$$

where $\mathcal{R} \equiv \mathcal{R}^{(1)}\otimes\mathcal{R}^{(2)}$ (in a Sweedler notation with upper indices and suppressed summation index) denotes the quasitriangular structure of $U_qso(N)$. The action (which here and in [9] we choose to be *right*) on $\mathcal{DC}^*$ is completely specified by the transformation laws of the generators $\sigma^\alpha = x^\alpha, \xi^\alpha, \partial^\alpha$, which read

$$\sigma^\alpha \triangleleft g = \rho^\alpha_\beta(g)\sigma^\beta \qquad \sigma^\alpha = x^\alpha, \xi^\alpha, \partial^\alpha, \qquad g \in U_qso(N),$$
$$x^\alpha \triangleleft \eta = x^\alpha, \qquad \xi^\alpha \triangleleft \eta = \xi^\alpha, \qquad \partial^\alpha \triangleleft \eta = -\partial^\alpha;$$

here ρ denotes the N-dimensional representation of $U_qso(N)$. The elements

$$Z^\alpha_\beta := T^{(1)}\rho^\alpha_\beta(T^{(2)}), \qquad T := \mathcal{R}_{21}\mathcal{R} \equiv T^{(1)}\otimes T^{(2)}, \qquad \mathcal{R}_{21} \equiv \mathcal{R}^{(2)}\otimes\mathcal{R}^{(1)}$$

are generators of $U_qso(N)$, and make up the "$SO_q(N)$ vector field matrix" Z [17, 15]. We recall that the ribbon element $w \in U_qso(N)$ is the central element such that

$$w^2 = u_1 S(u_1), \qquad u_1 := (S\mathcal{R}^{(2)})\mathcal{R}^{(1)}.$$

It is well-known [5] that there exist isomorphisms $U_hso(N)[[h]] \simeq Uso(N))[[h]]$ of $\star$-algebras over $\mathbb{C}[[h]]$. This essentially means that it is possible to express the elements of either algebra as power series in $h = \ln q$ with coefficients in the other. In particular w has an extremely simple expression in terms of the quadratic Casimir C of $so(N)$:

$$w = q^{-C} = e^{-hC} = \mathbf{1} + O(h), \qquad C := X^a X_a =: L(L+N-2). \tag{25}$$

($\{X^a\}$ is a basis of $so(N)$). We denote by $\nu := w^{1/4}$ and by $\tilde{w}, \tilde{\nu}, \tilde{T}, \tilde{Z}$ the analogs of w, ν, T, Z obtained by replacing $\mathcal{R}$ by $\tilde{\mathcal{R}}$. As an immediate consequence

$$\tilde{w} = q^{-C} q^{-\eta^2}, \qquad \tilde{\nu} = q^{-C/4} q^{-\eta^2/4}, \qquad \tilde{T} = T q^{2\eta\otimes\eta}.$$

As shown in [7, 3, 10], for real q there exists a $\star$-algebra homomorphism[3]

$$\varphi : \mathcal{H} \bowtie \widetilde{U_q so(N)} \to \mathcal{H}, \tag{26}$$

acting as the identity on $\mathcal{H}$ itself,

$$\varphi(a) = a \qquad\qquad a \in \mathcal{H}. \tag{27}$$

(This requires introducing an additional generator $\eta' = \varphi(\eta) \in \mathcal{DC}^*$ subject to the condition $\varphi(q^\eta) = q^{\eta'} = q^{-N/2}\Lambda$, so that $[\eta', \xi^\alpha] = 0$, $[\eta', x^\alpha] = -x^\alpha$, $[\eta', \partial^\alpha] = \partial^\alpha$. In the sequel we shall often use the the shorthand notation

$$\varphi(g) =: g', \qquad\qquad g \in \widetilde{U_q so(N)}.$$

One finds [9] on the spherical harmonics of level l (with $l = 0, 1, 2, ...$)

$$\nu' S_l^I = q^{-l(l+N-2)/4}\, S_l^I = S_l^I \triangleleft \nu. \tag{28}$$

$\star$-Structure and Hermiticity in configuration space representation

Theorem 2.1 *[9] For $q > 0$ the $\star$-structure of $\mathcal{DC}^*$ given in [6, 14] can be expressed in the form*

$$x^{\alpha\star} = x^\alpha, \tag{29}$$

$$\xi^{\alpha\star} = q^N\, \xi^\beta\, Z'^{\alpha}_{\beta} \Lambda^{-2}, \tag{30}$$

$$\partial^{\alpha\star} = -\tilde{\nu}'^{-2} \partial^\alpha \tilde{\nu}'^2 = -q^{\frac{1-N}{2}}\, \tilde{\nu}'^{-2} \partial_\alpha \tilde{\nu}'^2 \Lambda, \tag{31}$$

$$d^\star = -\tilde{\nu}'^2 d \tilde{\nu}'^{-2}. \tag{32}$$

---

[3] This is the q-deformed analog of the realization in the $q = 1$ case of the Cartan-Weyl $L^{\alpha\beta}$ generators of $\widetilde{Uso(N)}$ through vector fields (i.e. 1st-order differential operators):

$$\varphi(L^{\alpha\beta}) = x^\alpha \partial^\beta - x^\beta \partial^\alpha, \qquad \varphi(x^\alpha) = x^\alpha \qquad \varphi(\partial_\alpha) = \partial_\alpha, \qquad \varphi(\eta) = -x^\alpha \partial_\alpha.$$

Integration over $\mathbb{R}_q^N$.

Up to normalization this is defined by [16]

$$\int_q f(x)\,d^Nx=\int\limits_0^\infty dr\,m(r)\,r^{N-1}\int_{S_q^{N-1}} d^{N-1}t\,f(t,r)=\int\limits_0^\infty dr\,m(r)\,r^{N-1}f_0(r). \tag{33}$$

where f_0 denotes the $l=0$ component in (13). The'weight' m along the radial direction r has to fulfill, beside $m(r)>0$, the q-scaling condition

$$m(r)=m(qr). \tag{34}$$

Among the weights fulfilling the latter we can single out two "extreme" cases:

1. $m(r)\equiv 1$;
2. $m(r)\equiv m_{J,r_0}(r):=|q-1|\sum\limits_{n=-\infty}^{\infty} r\delta(r-r_0q^n)$ ('Jackson' integral).

Integration over $\mathbb{R}_q^N$ fulfills the standard properties of reality, positivity, $\widetilde{U_qso(N)}$-invariance, and a slightly modified cyclicity property [16]. Moreover, if f is a regular function decreasing faster than $1/r^{N-1}$ as $r\to\infty$ the Stokes theorem holds

$$\int_q \partial_\alpha f(x)\,d^Nx=0, \qquad\qquad \int_q \hat{\partial}_\alpha f(x)\,d^Nx=0. \tag{35}$$

Integration of functions immediately leads to integration of N-forms ω_N: upon moving all the ξ's to the right of the x's we can express ω_N in the form $\omega_N=fd^Nx$ (d^Nx denotes the unique independent exterior N-form), and just have to set

$$\int_q \omega_N=\int_q f\,d^Nx. \tag{36}$$

We introduce the scalar product of two "wave-functions" $\boldsymbol{\phi},\boldsymbol{\psi}\in F$ and more generally of two "wave-forms" $\alpha_p,\beta_p\in\Omega^p$ by

$$(\boldsymbol{\phi},\boldsymbol{\psi}):=\int_q \boldsymbol{\phi}^\star\tilde{\nu}'^{-2}\boldsymbol{\psi}\,d^Nx, \qquad\qquad (\alpha_p,\beta_p):=\int_q \alpha_p^\star\ {}^*\ \tilde{\nu}'^{-2}\beta_p. \tag{37}$$

Setting $U^{-1}{}^{\delta}_{\gamma}:=g^{\lambda\delta}g_{\lambda\gamma}$, the second can be easily expressed in terms of integrals

$$(\alpha_p,\beta_p)=\frac{1}{c_{N-p}}\int_q \alpha^\star_{\gamma_p\ldots\gamma_1}U^{-1}{}^{\delta_p}_{\gamma_p}\ldots U^{-1}{}^{\delta_1}_{\gamma_1}\tilde{\nu}'^{-2}\beta_{\delta_p\ldots\delta_1}\,d^Nx \tag{38}$$

involving the form components of the generic $\omega_p\in\Omega^p$ in the frame basis:

$$\omega_p=\theta^{\gamma_1}\ldots\theta^{\gamma_p}\omega_{\gamma_p\ldots\gamma_1}.$$

[Note that for $q \neq 1$ $\omega_{\gamma_p \ldots \gamma_1} \in \mathcal{H} \setminus F$, because $\theta^a \in \mathcal{DC}^* \setminus \Omega^*$. As an integrand funtion at the rhs(38) we mean what one obtains after letting all ∂_α present under the integral sign act on functions on their right, using the derivation rule (7)].

As a consequence of Theorem 2.1 we obtain

$$(\alpha_p, \hat{d}\beta_{p-1}) = (\hat{\delta}\alpha_p, \beta_{p-1}), \qquad\qquad (\hat{d}\beta_{p-1}, \alpha_p) = (\beta_{p-1}, \hat{\delta}\alpha_p). \tag{39}$$

and the (formal) hermiticity of both the momenta $p^\alpha = i\partial^\alpha$ and the Laplacian Δ:

$$(\boldsymbol{\phi}, p^\alpha \boldsymbol{\psi}) = (p^\alpha \boldsymbol{\phi}, \boldsymbol{\psi}), \qquad\qquad (\alpha_p, \Delta\beta_p) = (\Delta\alpha_p, \beta_p). \tag{40}$$

The kinetic term in the action for a (real) p-form (i.e. an antisymmetric tensor with p-indices) field theory of mass M can be now introduced as

$$\mathcal{S}_k = (\Delta\alpha_p, \alpha_p) + M^2(\alpha_p, \alpha_p). \tag{41}$$

Spectral analysis for p^α, Δ is affordable (see e.g. [8]), because of the relatively simple derivation rules (7), compared e.g. to those of $p^\alpha_R := p^\alpha + p^{\alpha\star}$.

The above considerations are still *formal* unless we make sense out of

$$\tilde{\nu}' = q^{-C'/4} q^{-\eta'^2/4}$$

and its inverse as *positive-definite* (pseudodifferential) operators on suitable Hilbert subspaces of F, Ω^*. Now $q^{-C'/4}$ is well-defined and positive-definite by (12), (28). To define the action of $q^{-\eta'^2/4}$ on functions $\phi(r)$ we change variables $r \to y = \ln r$, whereby $\eta' = -\partial_y - N/2$ and $r^{N-1}dr = e^{Ny}dy$, for any function $\phi(r)$ denote $\tilde{\phi}(y) := \phi(r)$, and express $e^{yN/2}\tilde{\phi}(y) = r^{N/2}\phi(r)$ in terms of its Fourier transform,

$$e^{\frac{N}{2}y}\tilde{\phi}(y) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \hat{\phi}(\omega) e^{i\omega y} d\omega. \tag{42}$$

One finds that $q^{-\eta'^2/4}$ acts as multiplication by $q^{\omega^2/4}$ on the Fourier transform:

$$e^{\frac{N}{2}y} q^{-\eta'^2/4}\tilde{\phi}(y) = q^{-\partial_y^2/4} e^{\frac{N}{2}y}\tilde{\phi}(y)| = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} d\omega \hat{\phi}(\omega) e^{i\omega y} q^{\omega^2/4}.$$

For $q < 1$ the UV convergence of scalar products is drastically improved by $q^{\omega^2/4}$. Introducing tentatively the ‘Hilbert space of square integrable functions on $\mathbb{R}^N_q$’

$$L_2^m := \left\{ \mathbf{f}(x) \equiv \sum_{l=0}^{\infty} \sum_I S^I_l \, f_{l,I}(r) \in F \;\mid\; (\mathbf{f}, \mathbf{f}) < \infty \right\} \tag{43}$$

one finds [9] that all works automatically with $L_2^{m\equiv 1}$ (i.e. if we choose $m(r) \equiv 1$). Otherwise, we have to

1. further require that m is invariant under inversion, $m(r^{-1}) = m(r)$, and fulfills condition (A.26) in [9] in order that the scalar product (,) be positive-definite;
2. at rhs(43) replace F with the subspace $E^{m,\beta,n}$ whose elements $f_{l,I}(r)$'s admit each an analytic continuation in the complex r-plane with poles possibly only in

$$r_{j,k} := q^{j+\beta} e^{i\frac{\pi(2k+1)}{n}}, \tag{44}$$

where $\beta \in \{0, 1/2\}$, n is an integer submultiple of N, $k = 0, 1, ..., n-1$, and $j \in \mathbb{Z}$, in order that the hermiticity (40) is actually implemented.

Condition 1. is fulfilled by a large class of weights, including the 'Jackson' ones m_{J,q^β}. Condition 2. selects interesting subclasses of functions, including q-special functions with "quantized parameters", among the simplest one e.g. $1/[1+(q^j r)^n]$.

References

[1] U. Carow-Watamura, M. Schlieker, S. Watamura, *$SO_q(N)$ covariant differential calculus on quantum space and quantum deformation of Schrödinger equation*, Z. Phys. C **49** (1991), 439.

[2] B. L. Cerchiai, G. Fiore, J. Madore, *Geometrical Tools for Quantum Euclidean Spaces*, Commun. Math. Phys. **217** (2001), 521-554. math.QA/0002007

[3] C.-S. Chu, B. Zumino, *Realization of vector fields for quantum groups as pseudodifferential operators on quantum spaces*, Proc. XX ICGTMP, Toyonaka 1995; q-alg/9502005.

[4] A. Dimakis and J. Madore, *Differential calculi and linear connections,* J. Math. Phys. **37** (1996), 4647-4661.

[5] V. G. Drinfeld, *Quasi Hopf Algebras*, Leningrad Math. J. **1** (1990), 1419.

[6] L. D. Faddeev, N. Y. Reshetikhin, L. Takhtadjan, *Quantization of Lie groups and Lie algebras*, Leningrad Math. J. **1** (1990), 193-225.

[7] G. Fiore, *Realization of $U_q(so(N))$ within the Differential Algebra on* $\mathbb{R}_q^N$, Commun. Math. Phys. **169** (1995), 475-500.

[8] G. Fiore, *The Euclidean Hopf algebra $U_q(e^N)$ and its fundamental Hilbert space representations*, J. Math. Phys. **36** (1995), 4363-4405.

[9] G. Fiore, *On the hermiticity of q-differential operators and forms on the quantum Euclidean spaces* $\mathbb{R}_q^N$, math.QA/0403463

[10] G. Fiore, *Quantum group covariant (anti)symmetrizers, ε-tensors, vielbein, Hodge map and Laplacian*, J. Phys. A: Math Gen. **37** (2004), 9175-9193; and references therein.

[11] S. Majid, *Braided Momentum Structure of the q-Poincare Group*, J. Math. Phys. **36** (1995), 1991-2007.

[12] T. Masuda, Y. Nakagami, S. L. Woronowicz, *A C^* algebraic framework for quantum groups*, math.QA/0309338.

[13] O. Ogievetsky *Differential operators on quantum spaces for $GL_q(n)$ and $SO_q(n)$*, Lett. Math. Phys. **24** (1992), 245.

[14] O. Ogievetsky, B. Zumino *Reality in the Differential calculus on the q-Euclidean Spaces*, Lett. Math. Phys. **25** (1992), 121-130.

[15] P. Schupp, P. Watts, and B. Zumino, *Bicovariant Quantum Algebras and Quantum Lie Algebras*, Commun. Math. Phys. **157** (1993), 305; and references therein.

[16] H. Steinacker, *Integration on quantum Euclidean space and sphere in N dimensions*, J. Math Phys. **37** (1996), 4738.

[17] B. Zumino, *Introduction to the Differential Geometry of Quantum Groups*, In: K. Schmüdgen (ed.), Math. Phys. X, Proc. X-th IAMP Conf. Leipzig (1991), Springer-Verlag (1992).